

Computer and Computing Technologies in Agriculture II, Volume 2

Edited by
Daoliang Li
Chunjiang Zhao



Springer



ifip

METHOD FOR ACHIEVING IRREGULAR BOUNDARY AREA FOR COMPLETE FLUIDIC SPRINKLER	901
<i>Junping Liu, Shouqi Yuan, Hong Li, Xingye Zhu</i>	
IRRIGATION UNIFORMITY WITH COMPLETE FLUIDIC SPRINKLER IN NO-WIND CONDITIONS	909
<i>Xingye Zhu, Shouqi Yuan, Hong Li, Junping Liu</i>	
STRUCTURAL BREAKS AND THE RELATIONSHIP BETWEEN SOYBEAN AND CORN FUTURES PRICES ON THE DALIAN COMMODITY EXCHANGE OF CHINA	919
<i>Rufang Wang, Yonghong Du, Jian Wang</i>	
DESIGN OF MEASURE AND CONTROL SYSTEM FOR PRECISION PESTICIDE DEPLOYING DYNAMIC SIMULATING DEVICE	927
<i>Yong Liang, Pingzeng Liu, Lu Wang, Jiping Liu, Lang Wang, Lei Han, Xinxin Yang</i>	
THE CROP DISEASE AND PEST WARNING AND PREDICTION SYSTEM	937
<i>Juhua Luo, Wenjiang Huang, Jihua Wang, Chaoling Wei</i>	
KNOWLEDGE ACQUISITION AND REPRESENTATION OF THE GENERAL AUXILIARY DIAGNOSIS SYSTEM FOR COMMON DISEASE OF ANIMAL	947
<i>Jianhua Xiao, Hongbin Wang, Ru Zhang, Peixian Luan, Lin Li, Danning Xu</i>	
THE DEVELOPMENT OF A GENERAL AUXILIARY DIAGNOSIS SYSTEM FOR COMMON DISEASE OF ANIMAL	953
<i>Jianhua Xiao, Hongbin Wang, Ru Zhang, Peixian Luan, Lin Li, Danning Xu</i>	

It was displayed that the emergence grade of the wheat stripe rust was five and occurrence degree was outbreak in Xifeng region of Gansu province. The fact that the warning results coincided well with report results from plant protecting station of Gansu province showed that this warning system and prediction method were all feasible.

4. RESULT

A warning and information system for diseases and pests of crops was established base on Geography Information System (GIS). The occurrence and damage of major diseases and pests of crops could be monitored and forecasted in real time by using the system, based on the forecasting model. And the system could submit warning maps with 6 colors and words information to the users. By using the system, we could standardize the forecasting data collection, transmit information through network and forecasting results viewable. Users could use it to classify and predict diseases and pests, also could use it to select fitting time and technology to control.

5. DISCUSSION

1) With the development of all kinds of subjects and lots of technology, many warning systems had been established at home and abroad. However, the system which was based on GIS was different from others traditional prediction system. The reasons were as follows: first, this system was based on .NET platform, and the interface was friendly, so the system was stable, portable, extensible and flexible. Second, this system could not only realize single point prediction but also regional prediction by basing on the function of GIS spatial analysis and multiple spatial interpolation methods. Finally, the remote sense was developed in the warning system to obtain fast and real-time the information of diseases and insects, which improved the accuracy of early warning result to a certain extent (BONE, S et al,2005).

2) The models in the warning system were all from literature data, so the models could only be used in some regions, which might result in the deviation of warning between theoretical value and actual value. However, with development and wide application of other information technology, the subsystem of automatic modeling could be established and used in the warning system to improve the accuracy of prediction.

THE CROP DISEASE AND PEST WARNING AND PREDICTION SYSTEM

Juhua Luo^{1,2}, Wenjiang Huang^{1*}, Jihua Wang¹, Chaoling Wei²

¹ National Engineering Research Center For Information Technology In Agriculture, Beijing, P. R. China 100097

² Resources and Environment College of Anhui Agricultural University, Hefei, Anhui Province, P. R. China 230036

*Corresponding author, Address: National Engineering Research Center for Information Technology In Agriculture, Beijing 100097, P. R. China, Tel: +86-10-51503647, Fax: +86-10-51503750, Email: yellowstar0618@163.com

Abstract: The aim of this study was to establish the warning and prediction system for crop diseases and pests based on SuperMap IS. NET geographic information system (GIS), which was developed by Supermap company. In this system, the author used GIS and remote sensing (RS) technology. The system could transform data information into a geographical information map to show the occurrence degree and distribution on various diseases and pests. This paper described mainly warning flow, database design and the main functions of the system. Finally, the system realized successfully the warning of the wheat stripe rust in Xifeng region of Qingyang city in Gansu province in 2002, and the prediction result was satisfactory. It indicated that we could classify and predict diseases and pests, and select right time and technology to control the diseases and pests by this GIS system.

Key words: disease and pest; geographic information system (GIS); Warning and prediction system

1. INTRODUCTION

The crop diseases and insect pests were all important biological hazard in agricultural production, and for many years, they restricted seriously the agricultural sustainable development. Statistics from the UN Food and

Please use the following format when citing this chapter:

Luo, J., Huang, W., Wang, J. and Wei, C., 2009, in IFIP International Federation for Information Processing, Volume 294, *Computer and Computing Technologies in Agriculture II, Volume 2*, eds. D. Li, Z. Chunjiang, (Boston: Springer), pp. 937-945.

3) Some standards and measures could be made for the investigation and controlling diseases and pests according to the pre-warning results of the warning system.

2.2 The design and realization the warning system of crops diseases and pests

2.2.1 Design route

At first, it was important to choose feasible GIS platform and the prediction model on diseases and pests after requirement analysis. Second, database was established according to parameter of model, and function modules of the system were also designed. Finally, the warning system was established and published after debugging the system (Feng Jiangfan et al,2006).

2.2.2 Platform selection

SuperMap IS. NET was chosen as the system development platform, the reasons were as follows:

- 1) Components of design could be easily managed.
- 2) The Multi-source data could be integrated and the massive image could be quickly accessed to.
- 3) Server was clustering; it was with a high degree of flexibility
- 4) In the platform, Client and server belonged multi-level cache structure, which could support a variety of map engine work together.

2.2.3 Database design

1) Attribute database

The attribute database of the system was composed with monitoring data, basic data on diseases and pests, national meteorological observation data and the data table about latitude and longitude of meteorological observation site, and so on (Wang Minghong et al,2006).

Monitoring data on diseases and pests: field investigation data, monitoring data using remote sensing, field experiments data, and so on.

Basic data on diseases and pests: diseases and pests species data, damage characteristics data, control and preventive methods data, and so on.

National meteorological observation data: temperature, moisture, rainfall and sunshine number, and so on.

2) Spatial databases

The spatial database of the warning was composed of basic map, thematic

map and warning information map.

Basic map: national administrative division map, digital elevation map.

Thematic map was got by basic map according to some goal, which included weather map, remote sensing imagery and crop division map.

Warning information map was obtained by spatial analysis on the basis of few thematic maps and analysis result of prediction model, by which the system could show the occurrence and distribution on variety of diseases and pests. And ultimately, some effective guidance can be given to prevent and control the diseases and pests according the warning information map.

2.2.4 Basic functions of the system

The functions of warning system was composed of special functions and general functions which were offered by GIS platform, including the translating, mitigating and amplifying graphics, and adding and deleting map layer, and so on. While the special functions were as follows:

1) Retrieval of diseases and insect pests

Users could search all kinds of knowledge about diseases and pests, for example, latin name and English name of disease and pest, harm symptoms and epidemic law on diseases and pests and control methods, and so on (Zhou Qiang, et al, 2004).

2) Diagnosis of diseases and pests

Users could obtain some papers on warning results and prediction methods after the operations of systems by various data and prediction models of diseases and pests.

3) Model base management

Administrators could add, delete, or modify the models in the system by Model base management (Liu Shuhua et al, 2003).

4) Monitoring and evaluation using remote sensing

The function was realized by the module of pests monitoring. Multiple and multi-temporal remote data was applicated in the module, and assessment model based on RS was call in the module (SONG Y H et al, 1993). By the function module, users could develop monitoring and assessment the diseases and pests. And related remote sensing image s could also were call from spatial database and displayed for users.

5) Visualization of warning result

By making visualization and spatial analysis with GIS, the system could transform prediction result and emergence grade into a geographical information map to display visually the occurrence and distribution on variety of diseases and pests. And ultimately, some effective guidance could be given by the system to prevent and control the diseases and pests (Si Lili et al, 2006).

Warning flow

The warning system was established on the basis of a database which was composed of attribute database and spatial database and a model base. At first, the result of diseases models and pests prediction models was transformed into thematic maps from spatial database and was overlaid on the information map. Finally, the system could offer a timely electronic information map which could show the occurrence and distribution on variety of diseases and pests (Fig. 1).




Fig. 1. Flow chart of warning system for diseases and pests

THE APPLICATION INSTANCE OF THE WARNING SYSTEM

The process how to realize the function of the warning system was improved by taking the wheat stripe rust disease in Xifeng of Qingyang

region in Gansu province as an example.

3.1 Summary of study area

The wheat stripe rust disease was one of the most serious diseases in China. Xifeng of Qingyang region in Gansu province was taken as an example to display the process how to realize the function of the warning system.

It was reported that the occurrence of the wheat stripe rust disease was mildly severe and severe symptoms in 1986, 1989, 1990, 1991, 2002, 2003, outbreak in 1985, moderate in 1993, and mild or none occurrence in other years. The wheat stripe rust disease was intermittent disease, but the occurrence was continuous and frequent, and becoming more and more serious in Xifeng of Qingyang region in Gansu province in recent ten years.

3.2 Data sources

- 1) The data of wheat stripe rust disease was from plant protecting station in Qingyang region in Gansu province;
- 2) The weather data was offered by meteorological observing station in Qingyang region in Gansu province;

3.3 Model selection

At present, the models in the system were all from literature data. The model which was selected according to expert experience and model characteristics was as follows:

$$y = 0.3251 + 0.0414x_1 + 0.0243x_2 + 0.0667x_3$$

In above prediction model, y is emergence grade, x_1 is average temperature from January to March, x_2 is index of temperature and rainfall, x_3 is index of the disease.

3.4 Release the warning information according to the model

3.4.1 Grading standard on the stripe rust wheat

Six degrees was adopted to evaluate the emergence of the wheat stripe rust, none, mild, lower than moderate, moderate, more than moderate, outbreak and were correspondingly marked as 0, 1, 2, 3, 4, 5 and identified as white, blue, green, yellow, orange, red (table 1).

Table 1 Grading standards for occurrence degree of wheat stripe rust disease

emergence grade	occurrence degree	warning color
none	none	white
one	mild	blue
two	lower than moderate	green
three	moderate	yellow
four	more than moderate	orange
five	outbreak	red

3.4.2 The pre-warning results on study area

1) System interface

The left page of the warning system concluded the field of 'user' logging and operational area of prediction, while the data about the warning was display on the right of system interface with victor diagram.

2) The operating process of the warning on the wheat stripe rust disease

At first, the users could enter the main interface of the system by login; second, users could enter the main interface of diseases prediction by clicking the diseases prediction model. Finally, used could develop step-by-step operation in the operational area of prediction according.

3) The pre-warning results on study area

By the above operation, users could call, in the model from the database of the system, and got the warning result by calculation. Finally, the pre-warning results were displayed on the right of the main interface by different color by electrical information map, at the same time, the warning results were also display by character form on the left of main interface (Fig.2).



Fig.2 Pre-warning results of wheat stripe rust disease in Xifeng county of Qingyang city in 2002

ACKNOWLEDGEMENTS

This work was subsidized by the National High Tech R&D Program of China (2006AA10Z203, 2007AA10Z201), National Natural Science Foundation of China (40701119), and funds (2006BAD10A01, 2007BAH12502). The authors are grateful to Mr. Weiguo Li, and Mrs. Hong Chang for data collection.

REFERENCES

- C. BONE, S. DRAGICEVIC and A. ROBERTS. Integrating high resolution remote sensing, GIS and fuzzy set theory for identifying susceptibility areas of forest insect infestations[J]. *International Journal of Remote Sensing*. 2005,26(21):4809-4828
- Hong Jiangfan, Ying Xuewei, Zhang Hong, et al. Forewarning System of Blue Algae for Taihu Lake Based on GIS. *Environmental Science & Technology*. 2006,29(9):59-62
- Huang Mayi, Wang Jihua, et al. Hyperspectral character of stripe rust on winter wheat and monitoring by remote sensing. *Journal of Agricultural Engineering Research*. 2003,19(6):154-158
- Huang Jinbao, Chen Yunhao, Huang Wenjiang, et al. Hyperspectral estimation models for LTN content of winter wheat canopy under stripe rust stress. *Journal of Agricultural Engineering Research*. 2008,24(1):34-39
- Lian Shuhua, Yang Xiaohong, et al. Decision support system for crop diseases and insect pests prevention and control based on GIS. *Journal of Agricultural Engineering Research*. 2003,19(4):148-150
- SHI H, Cao Keqiang, et al. Establishment of a real-time monitoring and forecasting system on main crop diseases and pests of China based on GIS. *Journal of Plant Protection*. 2006,33(3):282-286
- KING Y H, HEONG K L. Use of geographical Information system in analyzing large area distribution and dispersal of rice insects in South Korea[J]. *Journal of Applied Entomology*. 1993,32(3):307-316
- Wang Minghong, Jin Xiaohua, Liu Qian. The establishment of a long-distance warning and information system for major diseases and pests of crops in Beijing and its application. *China Plant Protection*. 2006,7(26):5-10
- Wang Shuhai, et al. Establishment of space information database of prediction systems of *Dendrolimus* sp. Based on GIS. *Journal of Liaoning Forestry Science & Technology*. 2005,4(13):35-36
- Zeng Shimai. *Macro-phytopathology*. Chinese Agricultural Technology Publishing House. 2005:182-184
- Zhou Qiang, Zhang Runjie. Studies of WebGIS-Based Warning Information System of Planthoppers Disaster. *ACTA Universitatis Sunyatseni*. 2004,30(21):64-66